Seizure Emergency Care Plan and				
ARENT/GUARDIAN complete and sign the top portion of form.				
Child Name:	Birth date:			Diago
Parent/Guardian Contact:	Phone:			Place
Emergency Contact:	Phone:			child's photo here
School:	Grade: □ flashing lights □ illness □ hunger □ temperature □ Other:			
Triggers: □ tiredness □ flashing lights Seizure Aura (if any): Seizure history: □ Convulsive □ Foca				
Describe:				
Antiseizure Medication Taken at Home	Common side	e effects		
Other Seizure Treatments/Special Diet T				
give permission for school personnel to share this in ealth care provider. I assume full responsibility for p ly child.				
ARENT SIGNATURE DA	 TE	SCHOOL NURSE SIGN	ATURE DATE	
EALTH CARE PROVIDER to complete al	l items. SIGN an	d DATE completed for	n.	
IF YOU SEE THIS:		DO THIS:		
☐ Convulsive Generalized Tonic Clonic:		Time the seizure		
You will see loss of consciousness. Stiffening of		Keep calm. Provide	reassurance.	
Rhythmic jerking movements. Convulsive seizures may last 1-5		Protect head, keep airway clear, turn on side if possible.		
minutes. The child may have a warning (aura) before the seizure.		<ul><li>4. Do not place anything in mouth.</li><li>5. Call 911 if student is injured or has difficulty breathing.</li></ul>		
Sleepiness and confusion may occur after the se	eizure.		s injured or has difficulty breathi	ng.
		6. Call parent. 7. Stay with student ur	ntil recovered from seizure.	
			treatments as marked below	,
		o. Administer rescue	treatments as marked below	•
<ul> <li>Focal:</li> <li>These seizures may begin with an aura. They m</li> </ul>	av he nartly alert	<ol> <li>Time the seizure</li> <li>Gently guide child a</li> </ol>	way from danger	
or unconscious. You may see lip smacking, chev			nd reassure them until recovere	d from seizure
or picking at clothes. These seizures usually last			that is stopped by a touch or a	
		5. Call parent.		
		6. Administer rescue	treatments as marked below	
☐ <b>Absence</b> : You will see quick changes in alert eye flutter or small twitching. Usually last less that		]		
Rescue Treatments				
☐ Child has a VNS. Child/staff may swipe with a	ura. Staff may swip	e at onset of seizure and e	very 60 seconds until seizure st	tops.
Give rescue medications below if seize	ure does not stop w	ithinminutes.		
If seizure <u>lasts longer</u> than minutes administ	er:			
☐ Diastatmg rectally	☐ Midazo	lammg in the nose	☐ Clonazepammg in	the cheek
☐ Multistep seizure rescue plan – Ple	ase see attached le	tter for details.		
·		<del></del> -		
	dminister:			<del> </del>
☐ Diastatmg rectally	☐ Midazo	lammg in the nose	☐ Clonazepammg in	the cheek
$\square$ Multistep seizure rescue plan – Ple	ase see attached le	tter for details.		
If emergency medication is administered:	Call 911 immediat	elv or □ Call 911 if seizur	e does not stop within 5 minu	utes
Other:			<u> </u>	
lf un amaguna and madication is at a shoot	ما المانية ما المسيد			
If no emergency medication is at school Call family to bring medications to school o				
ccommodations: Always take seizure act				ield trips.
Close adult supervision when swimming or c				u po.
The state of the s				
IEALTH CARE PROVIDER SIGNATURE		D'C NAME DUC	NF/FAX DATI	 E